

Roll No.

(07/20-I)

5260

B. A./B. Sc. EXAMINATION

(Sixth Semester)

MATHEMATICS

BM-362

Linear Algebra

Time : Three Hours Maximum Marks : $\begin{cases} \text{B.Sc. : 40} \\ \text{B.A. : 26} \end{cases}$

Note : Attempt *Five* questions in all, selecting *one* question from each Unit. Q. No. 1 is compulsory. Marks outside the brackets are for B.Sc. students and marks within the brackets are for B.A. students.

(Compulsory Question)

1. (a) Define a Vector Space. 2(1½)

(1-03/21) B-5260

P.T.O.

- (b) Show that the set $\{1, i\}$ generates \mathbb{C} (set of complex numbers) over \mathbb{R} (set of real numbers). $2(1\frac{1}{2})$
- (c) Define singular and non-singular linear transformation. $2(1\frac{1}{2})$
- (d) Define orthogonal set in an inner product space. $2(1\frac{1}{2})$

Unit I

2. (a) Prove that union of two subspaces of a vector space, is a subspace if and only if one is contained in the other. $4(2\frac{1}{2})$
- (b) Show that the set $\{(1, 2, 3), (0, 1, 2), (0, 0, 1)\}$ generates the vector space $\mathbb{R}^3(\mathbb{R})$. $4(2\frac{1}{2})$
3. (a) If $V(F)$ is a finitely generated vector space, then prove that any maximal linearly independent subset of $V(F)$ is a basis of $V(F)$. $4(2\frac{1}{2})$

- (b) If W is a subspace of a vector space $V(F)$ and $u, u' \in V$, then $W + u = W + u'$ iff $u - u' \in W$. $4(2\frac{1}{2})$

Unit II

4. (a) Show that linear transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by :

$$T(x_1, x_2) = (x_1 \cos \theta + x_2 \sin \theta, -x_1 \sin \theta + x_2 \cos \theta)$$

is a vector space isomorphism. $4(2\frac{1}{2})$

- (b) Let $T : U \rightarrow V$ be a linear transformation, then $U/\ker T \cong T(U)$. $4(2\frac{1}{2})$
5. (a) If $T : U \rightarrow V$ is a linear transformation, then $\text{Rank } T + \text{Nullity } T = \text{Dim. } U$. $4(2\frac{1}{2})$
- (b) Let $S = \{v_1, v_2, v_3\}$ be a basis of $V_3(\mathbb{R})$, defined by $v_1 = (-1, 1, 1)$, $v_2 = \{1, -1, 1\}$, $v_3 = \{1, 1, -1\}$. Find the dual basis of S . $4(2\frac{1}{2})$

Unit III

6. (a) Let $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be a linear operator defined by $T(x, y, z) = (2x, 4x - y, 2x + 3y - z)$. Show that T is invertible and find T^{-1} . $4(2\frac{1}{2})$
- (b) If $C(\mathbb{R})$ is a vector space and T is a linear transformation on $C(\mathbb{R})$ defined by $T(a + ib) = a - ib$ for all $a, b \in \mathbb{R}$; find the matrix of T with respect to ordered basis $B = \{1 + i, 1 + 2i\}$. $4(2\frac{1}{2})$
7. (a) Let U and V be finite dimensional vector spaces and let $B = \{u_1, u_2, \dots, u_n\}$ and $B' = \{v_1, v_2, \dots, v_n\}$ be ordered basis for U and V respectively. If T is a linear transformation from U to V then for any $u \in U$:

$$[T(u), B'] = [T : B, B'] [u, B].$$

$4(2\frac{1}{2})$

- (b) Show that the matrix $A = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}$ is not diagonalizable over the field \mathbb{C} . $4(2\frac{1}{2})$

Unit IV

8. (a) State and prove Cauchy Schwarz inequality. $4(2\frac{1}{2})$
- (b) Prove that every finite dimensional vector space is an inner product space. $4(2\frac{1}{2})$
9. (a) Obtain an orthonormal basis with respect to standard inner product for subspace of \mathbb{R}^3 generated by $(1, 0, 1)$, $(1, 0, -1)$ and $(0, 3, 4)$. $4(2\frac{1}{2})$
- (b) Let T be a linear operator on a finite dimensional inner product space $V(F)$. Show that T can be uniquely expressed as the sum of a self-adjoint and skew-symmetric operator. $4(2\frac{1}{2})$